

Fibre bundle formulation of relativistic quantum mechanics

Bozhidar Z. Iliev * † ‡

Short title: Bundle relativistic quantum mechanics

Basic ideas: → November 1997, January 1998

Began: → February 16, 1998

Ended: → March 27, 1998

Initial typeset: → April 1–14, 1998

Revised: → August 1999, March 2002

Last updated: → March 30, 2002

Produced: → February 1, 2008

LANL arXive server E-print No.: quant-ph/0201085

BOZHO[®] TM

Subject Classes:

Relativistic quantum mechanics, Differential geometry

2000 MSC numbers:

81Q99, 81S99

2001 PACS numbers:

02.40.Ma, 02.40.Yy

02.90.+p, 03.65.Pm

Key- Words:

*Relativistic quantum mechanics, Fibre bundles,
Geometrization of relativistic quantum mechanics,
Relativistic wave equations, Dirac equation, Klein-Gordon equation*

*Department Mathematical Modeling, Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Boul. Tzarigradsko chaussée 72, 1784 Sofia, Bulgaria

†E-mail address: bozho@inrne.bas.bg

‡URL: <http://theo.inrne.bas.bg/~bozho/>

Contents

1	Introduction	1
2	Bundle nonrelativistic quantum mechanics (review)	2
3	Time-dependent (Hamiltonian) approach	4
3.1	General case	4
3.2	Dirac equation	6
3.3	Klein-Gordon equation	7
3.4	Other relativistic wave equations	8
4	Covariant approach	9
4.1	Dirac equation	9
4.2	Other relativistic wave equations	12
5	Propagators and evolution transports or operators	13
5.1	Green functions (review)	13
5.2	Nonrelativistic case (Schrödinger equation)	14
5.3	Dirac equation	16
5.4	Klein-Gordon equation	17
6	Conclusion	18
	Appendix A Linear transports along maps in fibre bundles	18
	Appendix B Matrix operators	20
	References	22
	This article ends at page	24

Distribution of the material:

- I (1–3): Time-dependent approach, arXiv E-print No. quant-ph/0105056
 II (4&5 & A&B): Covariant approach, arXiv E-print No. quant-ph/0107002

Abstract

We propose a fibre bundle formulation of the mathematical base of relativistic quantum mechanics. At the present stage the bundle form of the theory is equivalent to its conventional one, but it admits new types of generalizations in different directions. In the bundle description the wavefunctions are replaced with (state) sections (covariant approach) or liftings of paths (equivalently: sections along paths) (time-dependent approach) of a suitably chosen vector bundle over space-time whose (standard) fibre is the space of the wavefunctions. Now the quantum evolution is described as a linear transportation of the state sections/liftings in the (total) bundle space. The equations of these transportations turn to be the bundle versions of the corresponding relativistic wave equations. Connections between the (retarded) Green functions of these equations and the evolution operators and transports are found. Especially the Dirac and Klein-Gordon equations are considered.

1. Introduction

This investigation is devoted to a pure geometric (re)formulation of the (mathematical) base of relativistic quantum mechanics in terms of fibre bundles. It can be regarded as a direct continuation of [1–7] where a full self-consistent fibre bundle version of non-relativistic quantum mechanics is elaborated. Many ideas of these works can be transferred to the relativistic case but, as we shall see below, they are not enough for the above aim. For example, the right mathematical concept reflecting the peculiarities of relativistic quantum evolution is the (linear) transport along the identity map of the space-time, on the contrary to a (linear) transport along paths (in space-time) in the non-relativistic case. A new problem is the fact that some relativistic wave equations are linear partial differential equations of second order; so for them the methods developed for the treatment of Schrödinger equation can not be applied directly.

The presented in [2, sec. 4.1] motivations for applying the theory of fibre bundles to non-relativistic quantum mechanics can *mutatis mutandis* be transferred in the case of relativistic quantum mechanics. For the purpose, first of all, one has to replace the 3-dimensional coordinate (or other) space model with the Minkowski spacetime. More specific changes are connected with particular relativistic wave equations one investigates.

In the bundle description the relativistic wavefunctions are replaced by (state) liftings of paths, sections or sections along paths of a suitable vector bundle with the space-time as a base and whose (standard, typical) fibre is the space where the corresponding conventional wavefunctions (or some combinations of them and their partial derivatives) ‘live’. The operators acting on this space are replaced with appropriate liftings of paths or morphisms along paths and the evolution of the wavefunctions is now described as a linear transport of the state liftings/sections (possibly along paths) in this bundle.

The lay-out of the paper is the following.

A review of the bundle approach to non-relativistic quantum mechanics is presented in Sect. 2. It contains certain basic ideas and equations of [2–7] required as a starting point for the present work.

Sect. 3 is devoted to the time-dependent or Hamiltonian bundle description of relativistic wave equations. This approach is a straightforward generalization of the bundle description of nonrelativistic quantum mechanics to relativistic one. In Subsect. 3.1 is developed a general scheme for fibre bundle treatment of Schrödinger-type partial differential equations. The method is also applicable to linear partial differential equations of higher orders. This is achieved by transforming such an equation to a system of first-order linear partial differential equations (with respect to a new function) which, when written in a matrix form, is just the Schrödinger-like presentation of the initial equation. Of course, this procedure is not unique and its concrete realization depends on the physical problem under exploration. In Subsect. 3.2 is given a bundle description of the Dirac equation. Now the corresponding vector bundle over the space-time has as a (standard) fibre the space of 4-spinors and may be called 4-spinor bundle. Since the Dirac equation is of first order, it can be rewritten in Schrödinger-type form. By this reason the formalism outlined in Sect. 2 is applied to it *mutatis mutandis*. In particular, here the state of a Dirac particle is described by a lifting of paths or section along paths of the 4-spinor bundle and equivalently rewrite Dirac equation as an equation for linear transportation of this section with respect to the corresponding *Dirac evolution transport* in this bundle. Subsect. 3.3 contains several procedures for transforming Klein-Gordon equation to Schrödinger-like one. After such a presentation is chosen, we can, analogously to Dirac case in Subsect. 3.2, apply to it the methods of Sect. 2 and Subsect. 3.1. Comments on fibre bundle description of other relativistic wave equations are given in Subsect. 3.4.

With Sect. 4 we begin to develop the covariant approach to bundle description of rel-

ativistic quantum mechanics. Subsect. 4.1 contains a covariant application of the ideas of Sect. 2 to Dirac equation. The bundle, where Dirac particles ‘live’, is a vector bundle over space-time with the space of 4-spinors as a fibre; so here we work again with the 4-spinor bundle of Subsect. 3.2, but now the evolution of a Dirac particle is described via a *geometric transport* which is a *linear transport along the identity map of space-time*. The state of a Dirac particle is represented by a section (not along paths!) of the 4-spinor bundle and is (linearly) transported by means of the transport mentioned. The Dirac equation itself is transformed into a covariant Schrödinger-like equation. In Subsect. 4.2 we apply the covariant bundle approach to Klein-Gordon equation. For this purpose we present a 5-dimensional representation of this equation as a first-order Dirac-like equation to which the theory of Subsect. 4.1 can be transferred *mutatis mutandis*.

The goal of Sect. 5 is to be revealed some connections between the retarded Green functions ($\equiv$ propagators) of the relativistic wave equations and the corresponding to them evolution operators and transports. Generally speaking, the evolution operators (resp. transports) admit representation as integral operators, the kernel of which is connected in a simple manner with the retarded Green function (resp. Green morphism of a bundle). Subsect. 5.1 contains a brief general consideration of the Green functions and their connection with the evolution transports, if any. In Subsect. 5.2, 5.3, and 5.4 we derive the relations mentions for Schrödinger, Dirac, and Klein-Gordon equations, respectively.

Sect. 6 closes the paper with a brief summary of the main ideas underlying the bundle description of relativistic quantum mechanics.

Appendix A contains some mathematical results concerning the theory of (linear) transports along maps required for the present investigation.

In Appendix B are given certain formulae concerning matrix operators, i.e. matrices with operator entries, which arise naturally in relativistic quantum mechanics.

2. Bundle nonrelativistic quantum mechanics (review)

In the series of papers [2–7] we have reformulated nonrelativistic quantum mechanics in terms of fibre bundles. The mathematical base for this was the Schrödinger equation

$$i\hbar \frac{d\psi(t)}{dt} = \mathcal{H}(t)\psi(t), \quad (2.1)$$

where $i \in \mathbb{C}$ is the imaginary unit, $\hbar (= h/2\pi)$ is the Planck constant divided by 2π , ψ is the system’s state vector belonging to an appropriate Hilbert space $\mathcal{F}$, and $\mathcal{H}$ is the system’s Hamiltonian. Here t is the time considered as an independent variable (or parameter). If ψ is known for some initial moment t_0 , the solution of 2.1 can be written as

$$\psi(t) = \mathcal{U}(t, t_0)\psi(t_0) \quad (2.2)$$

where $\mathcal{U}$ is the *evolution operator* of the system [8, chapter IV, sect. 3.2] (for details see [2]).

In the bundle approach the system’s Hilbert space $\mathcal{F}$ is replace with a Hilbert bundle (F, π, M) with (total, fibre) bundle space F , projection π , base M , isomorphic fibres $F_x := \pi^{-1}(x)$, $x \in M$, and (standard, typical) fibre coinciding with $\mathcal{F}$. So, there exist isomorphisms $l_x: F_x \rightarrow \mathcal{F}$, $x \in M$. In the present work the base M will be identified with the Minkowski space-time M^4 of special relativity.¹

¹The bundle formulation of nonrelativistic quantum mechanics is insensitive to the/this choice of M [6]; in it is more natural to identify M with the 3-dimensional Newtonian space $\mathbb{E}^3$ of classical mechanics.

In the *Hilbert bundle description* a state vector ψ and the Hamiltonian $\mathcal{H}$ are replaced respectively by a state section along paths $\Psi: \gamma \rightarrow \Psi_\gamma$ and the *bundle Hamiltonian* (morphism along paths) $H: \gamma \rightarrow H_\gamma$, given by [2, 3, 6]:

$$\Psi_\gamma: t \rightarrow \Psi_\gamma(t) = l_{\gamma(t)}^{-1}(\psi(t)), \quad H_\gamma: t \rightarrow H_\gamma(t) = l_{\gamma(t)}^{-1} \circ \mathcal{H}(t) \circ l_{\gamma(t)} \quad (2.3)$$

where $\gamma: J \rightarrow M$, J being an $\mathbb{R}$ -interval, is the world line (path) of some (point-like) observer, $t \in J$, and $\circ$ denotes composition of maps.

The bundle analogue of the evolution operator $\mathcal{U}$ is the *evolution transport* U along paths, both connected by

$$U_\gamma(t, s) = l_{\gamma(t)}^{-1} \circ \mathcal{U}(t, s) \circ l_{\gamma(s)}: F_{\gamma(s)} \rightarrow F_{\gamma(t)}, \quad s, t \in J, \quad (2.4)$$

which governs the evolution of state liftings via (cf. (2.2))

$$\Psi_\gamma(t) = U_\gamma(t, s)\Psi_\gamma(s), \quad s, t \in J. \quad (2.5)$$

The bundle version of (2.1), the so-called *bundle Schrödinger equation*, is [3, 7]

$$D\Psi = 0. \quad (2.6)$$

Here D is the derivation along paths corresponding to U , viz (cf. (A.9) and [9, definition 4.1]; see also [10] and [2, definition 3.4])

$$D: \text{PLift}^1(E, \pi, B) \rightarrow \text{PLift}^0(E, \pi, B)$$

where $\text{PLift}^k(F, \pi, M)$ is the set of C^k liftings of paths from M to F , and its action on a lifting $\lambda \in \text{PLift}^1(F, \pi, M)$ with $\lambda: \gamma \mapsto \lambda_\gamma$ is given via

$$D_s^\gamma \lambda := \lim_{\varepsilon \rightarrow 0} \left\{ \frac{1}{\varepsilon} [U_\gamma(s, s + \varepsilon)\lambda_\gamma(s + \varepsilon) - \lambda_\gamma(s)] \right\} \quad (2.7)$$

where $D_s^\gamma(\lambda) := ((D\lambda)(\gamma))(s) = (D\lambda)_\gamma(s)$.

If $\{e_a(\gamma(s))\}$, $s \in J$ is a basis in $F_{\gamma(s)}$, the explicit action of D is (cf. (A.11); see also [9, proposition 4.2] and [10])

$$D_s^\gamma \lambda = \left(\frac{d\lambda_\gamma^a(s)}{ds} + \Gamma_a^b(s; \gamma)\lambda_\gamma^b(s) \right) e_a(\gamma(s)). \quad (2.8)$$

Here the *coefficients* $\Gamma_a^b(s; \gamma)$ of U are defined by (cf. and (A.12))

$$\Gamma_a^b(s; \gamma) := \frac{\partial (U_\gamma(s, t))^b_a}{\partial t} \Big|_{t=s} = - \frac{\partial (U_\gamma(t, s))^b_a}{\partial t} \Big|_{t=s} \quad (2.9)$$

where $(U_\gamma(s, t))^b_a$ are given via $U(t, s)e_a(\gamma(s)) =: \sum_b (U_\gamma(s, t))^b_a e_b(\gamma(t))$ and are the local components of U in $\{e_a\}$.

There is a bijective correspondence between D and the (bundle) Hamiltonian expressed by²

$$\mathbf{\Gamma}_\gamma(t) := [\Gamma_a^b(t; \gamma)] = -\frac{1}{i\hbar} \mathbf{H}_\gamma^{\mathbf{m}}(t) \quad (2.10)$$

²We denote the matrix corresponding to some quantity, e.g. vector or operator, in a given field of bases with the same (kernel) symbol but in **boldface**; e.g. $\mathbf{U}_\gamma(t, s) := [(U_\gamma(s, t))^b_a]$.

with

$$\mathbf{H}_\gamma^{\mathbf{m}}(t) = i\hbar \frac{\partial \mathbf{U}_\gamma(t, t_0)}{\partial t} \mathbf{U}_\gamma^{-1}(t, t_0) = \frac{\partial \mathbf{U}_\gamma(t, t_0)}{\partial t} \mathbf{U}_\gamma(t_0, t).$$

being the *matrix-bundle Hamiltonian* (for details see [3])³

In the Hilbert space description of quantum mechanics to a dynamical variable $\mathbf{A}$ corresponds an observable $\mathcal{A}(t)$ which is a linear Hermitian operator in $\mathcal{F}$. In the Hilbert bundle description to $\mathbf{A}$ corresponds a Hermitian lifting A of paths whose restriction on $F_{\gamma(t)}$ is

$$A_\gamma(t) = l_{\gamma(t)}^{-1} \circ \mathcal{A}(t) \circ l_{\gamma(t)} : F_{\gamma(t)} \rightarrow F_{\gamma(t)}. \quad (2.11)$$

The mean value of $\mathbf{A}$ at a state characterized by a state vector ψ or, equivalently, by the corresponding to it state lifting Ψ_γ is

$$\langle \mathcal{A}(t) \rangle_\psi^t := \frac{\langle \psi(t) | \mathcal{A}(t) \psi(t) \rangle}{\langle \psi(t) | \psi(t) \rangle} = \langle A_\gamma(t) \rangle_{\Psi_\gamma}^t := \frac{\langle \Psi_\gamma(t) | A_\gamma(t) \Psi_\gamma(t) \rangle_{\gamma(t)}}{\langle \Psi_\gamma(t) | \Psi_\gamma(t) \rangle_{\gamma(t)}}. \quad (2.12)$$

Here

$$\langle \cdot | \cdot \rangle_x = \langle l_x \cdot | l_x \cdot \rangle, \quad x \in M \quad (2.13)$$

is the fibre Hermitian scalar product in (F, π, M) induced by the Hermitian scalar product $\langle \cdot | \cdot \rangle : \mathcal{F} \times \mathcal{F} \rightarrow \mathbb{C}$ in $\mathcal{F}$.

A summary of the above and other details concerning the Hilbert space and Hilbert bundle description of (nonrelativistic) quantum mechanics can be found in [6].

3. Time-dependent (Hamiltonian) approach

Consider now the pure mathematical aspects of the scheme described in Sect. 2. On one hand, it is essential to be notice that the Schrödinger equation (2.1) is a first order (with respect to the time⁴) linear partial differential equation solved with respect to the time derivative. On the other hand, the considerations of Sect. 2 are true for Hilbert spaces and bundles whose dimensionality is generically infinity, but, as one can easily verify, they hold also for spaces and bundles with finite dimension.

These observations, as we shall prove below, are enough to transfer the bundle nonrelativistic formalism to the relativistic region. We call the result of this procedure *time-dependent or Hamiltonian approach* as in it the time plays a privileged rôle and the relativistic covariance is implicit.

3.1. General case

Taking into account the above, we can make the following conclusion. Given a linear (vector) space $\mathcal{F}$ of C^1 functions $\psi : J \rightarrow \mathbb{C}$ with J being an $\mathbb{R}$ -interval. (We do not make any assumptions on the dimensionality of $\mathcal{F}$; it can be finite as well as countable or uncountable infinity.) Let $\mathcal{H}(t) : \mathcal{F} \rightarrow \mathcal{F}$, $t \in J$ be (possibly depending on t) linear operator. Consider the equation⁵

$$i\hbar \frac{\partial \psi(t)}{\partial t} = \mathcal{H}(t) \psi(t). \quad (3.1)$$

³Note, the constant $i\hbar$ in (2.10) comes from the same constant in (2.1).

⁴The dependence of ψ and $\mathcal{H}$ on the spatial coordinates (and momentum operators) is inessential for the present section and, respectively, is not written explicitly.

⁵We introduce the multiplier $i\hbar \neq 0$ from purely physical reasons and to be able to apply the results already obtained directly, without any changes.

For it are valid all of the results of Sect. 2, viz., for instance, there can be introduced the bundle (F, π, M) , the evolution operator $\mathcal{U}$, etc.; the relations between them being the same as in Sect. 2. If the vector space $\mathcal{F}$ is endowed with a scalar (inner) product $\langle \cdot | \cdot \rangle: \mathcal{F} \times \mathcal{F} \rightarrow \mathbb{C}$, then (2.13) induces analogous product in the bundle (i.e. in the bundle's fibres). Consequently, imposing condition (2.12), we get (2.11). Further, step by step, one can derive all of the results of [2–7].

Further we shall need a slight, but very important generalization of the above. Let now $\mathcal{F}$ be a vector space of vector-valued C^1 functions $\psi: J \rightarrow V$ with V being a complex vector space and $\mathcal{H}(t): \mathcal{F} \rightarrow \mathcal{F}$; the case just considered corresponds to $V = \mathbb{C}$, i.e to $\dim_{\mathbb{C}} V = 1$. It is not difficult to verify that all of the above-said, corresponding to $\dim_{\mathbb{C}} V = 1$, is also *mutatis mutandis* valid for $\dim_{\mathbb{C}} V \geq 1$. Consequently, the fibre bundle reformulation of the solution of (3.1), the operators, and scalar product(s) in $\mathcal{F}$ can be carried out in the general case when $\psi: J \rightarrow V$ with $\dim_{\mathbb{C}} V \geq 1$.

Suppose now K^m is the vector space of C^m , $m \in \mathbb{N} \cup \{\infty\} \cup \{\omega\}$, vector-valued functions $\varphi: J \rightarrow W$ with W being a vector space. Let $\varphi \in K^m$ satisfies the equation

$$f\left(t, \varphi, \frac{\partial \varphi}{\partial t}, \dots, \frac{\partial^n \varphi}{\partial t^n}\right) = 0 \quad (3.2)$$

where $f: J \times K^m \times K^{m-1} \times \dots \times K^{m-n} \rightarrow K^{m-n}$, $n \in \mathbb{N}$, $n \leq m$ is a map (multi)linear in φ and its derivatives. We suppose (3.2) to be solvable with respect to the highest derivative of φ , i.e. (3.2) to be equivalent to⁶

$$\frac{\partial^n \varphi}{\partial t^n} = G\left(t, \varphi, \frac{\partial \varphi}{\partial t}, \dots, \frac{\partial^{n-1} \varphi}{\partial t^{n-1}}\right) \quad (3.3)$$

for some map $G: J \times K^m \times K^{m-1} \times \dots \times K^{m-n+1} \rightarrow K^{m-n}$, linear in φ and its derivatives.

The already developed fibre bundle formalism for the (Schrödinger-type) equation (3.1) can be transferred to (3.3). This can be done in a number of different ways. Below we shall realize the most natural way, but one has to keep in mind that for a concrete equation another method may turn to be more useful, especially from the view-point of possible physical applications (see below Subsect. 3.3).

Defining $\mathcal{F} := K^m \times K^{m-1} \times \dots \times K^{m-n+1}$ and putting

$$\psi := \left(\varphi, \frac{\partial \varphi}{\partial t}, \dots, \frac{\partial^{n-1} \varphi}{\partial t^{n-1}}\right)^{\top} \quad (3.4)$$

with $\top$ being the transposition sign, we can transform (3.3) in the form (3.1) with ‘Hamiltonian’

$$\mathcal{H}(t) = i\hbar \times \begin{pmatrix} 0 & \text{id}_{K^{m-1}} & 0 & \dots & 0 \\ 0 & 0 & \text{id}_{K^{m-2}} & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & \text{id}_{K^{m-n+1}} \\ f_0(t)\text{id}_{K^m} & f_1(t)\text{id}_{K^{m-1}} & f_2(t)\text{id}_{K^{m-2}} & \dots & f_{n-1}(t)\text{id}_{K^{m-n+1}} \end{pmatrix} : \mathcal{F} \rightarrow \mathcal{F} \quad (3.5)$$

⁶Requiring the equivalence of (3.2) and (3.3), we exclude the existence of sets on which the highest derivative of φ entering in (3.2) may be of order $k < n$. If we admit the existence of such sets, then on them we have to replace n in (3.3) with k . (Note k may be different for different such sets.) The below-described procedure can be modified to include this more general situation but, since we do not want to fill the presentation with complicated mathematical details, we are not going to do this here.

which is a linear matrix operator (see Appendix B), i.e. a matrix of linear operators. Here id_X is the identity map of a set X and $f_i: J \rightarrow \mathbb{C}$, $i = 0, \dots, n-1$ define the (multi)linear map $G: J \times \mathcal{F} \rightarrow K^{m-n}$ by

$$G\left(t, \varphi, \frac{\partial \varphi}{\partial t}, \dots, \frac{\partial^{n-1} \varphi}{\partial t^{n-1}}\right) = f_0(t)\varphi + \sum_{i=1}^{n-1} f_i(t) \frac{\partial^i \varphi}{\partial t^i}. \quad (3.6)$$

In this way we have proved that (3.2) can equivalently be rewritten as a Schrödinger-type equation (3.1) with ‘Hamiltonian’ given by (3.5). Such a transformation is not unique. For example, one can choose the components of ψ to be any n linearly independent linear combinations of φ , $\partial \varphi / \partial t$, $\dots$, $\partial^{n-1} \varphi / \partial t^{n-1}$; this will result only in another form of the matrix (3.5). In fact, if $A(t)$ is a nondegenerate matrix-valued function, the change

$$\psi(t) \mapsto \tilde{\psi}(t) = A(t)\psi(t) \quad (3.7)$$

leads to (3.1) with Hamiltonian

$$\tilde{\mathcal{H}}(t) = A(t)\mathcal{H}A^{-1}(t) + \frac{\partial A(t)}{\partial t}A^{-1}(t). \quad (3.8)$$

Now to the equation (3.1), with $\mathcal{H}$ given via (3.5), we can apply the already described procedure for reformulation in terms of bundles.

3.2. Dirac equation

The relativistic quantum mechanics of spin $\frac{1}{2}$ particle is described by the Dirac equation (see [11, chapter 2], [12, chapters 1–5], and [13, part V, chapter XX]). The wave function $\psi = (\psi_1, \psi_2, \psi_3, \psi_4)^\top$ of such a particle is a 4-dimensional (4-component) spinor satisfying this equation whose Schrödinger-type form⁷ is (see, e.g. [13, chapter XX, § 6, equation (36)] or [12, chapter 1, § 3, equation (1.14)])

$$i\hbar \frac{\partial \psi}{\partial t} = {}^D\mathcal{H}\psi \quad (3.9)$$

where ${}^D\mathcal{H}$ is a Hermitian operator, called *Dirac Hamiltonian*, in the space $\mathcal{F}$ of state vectors (spinors). For a spin $\frac{1}{2}$ particle with (proper) mass m and electric charge e the explicit form of ${}^D\mathcal{H}$ in an (external) electromagnetic field with 4-potential $(\varphi, \mathbf{A})$ is [13, chapter XX, § 6, equation (44)]

$${}^D\mathcal{H} = e\varphi \mathbb{1}_4 + c\boldsymbol{\alpha} \cdot (\mathbf{p} - \frac{e}{c}\mathbf{A}) + mc^2\beta, \quad (3.10)$$

where $\mathbb{1}_4$ is the 4×4 unit matrix, c is the velocity of light in vacuum, $\boldsymbol{\alpha} := (\alpha^1, \alpha^2, \alpha^3)$ is a matrix vector of 4×4 matrices, $\mathbf{p} = -i\hbar \nabla$ is the (3-dimensional) momentum operator ($\nabla_i := \partial / \partial x^i$, $i = 1, 2, 3$), and β is a 4×4 matrix. The explicit (and general) forms of $\alpha^1, \alpha^2, \alpha^3$, and β can be found, e.g. in [11, chapter 2, sect. 2.1.2, equation (2.10)].

Since (3.9) is a first-order equation, we can introduce the *Dirac evolution operator* ${}^D\mathcal{U}$ via $\psi(t) = {}^D\mathcal{U}(t, t_0)\psi(t_0)$ (cf. (2.2)). Generally it is a 4×4 integral-matrix operator (see Appendix B) uniquely defined by the initial-value problem

$$i\hbar \frac{\partial}{\partial t} {}^D\mathcal{U}(t, t_0) = {}^D\mathcal{H}(t) \circ {}^D\mathcal{U}(t, t_0), \quad {}^D\mathcal{U}(t_0, t_0) = \text{id}_{\mathcal{F}} \quad (3.11)$$

⁷For the purposes of this section we do not need the widely know relativistic invariant form of Dirac equation [11–13] (see also Subsect. 4.1).

with $\mathcal{F}$ being the space of 4-spinors. The explicit form of ${}^D\mathcal{U}$ is derived, e.g. in [11, chapter 2, sect. 2.5.1].

Now the bundle formalism developed in [2–7] can be applied to a description of Dirac particles practically without changes. For instance, the spinor lifting of paths have to be introduced via (2.3) and are connected by (2.5) in which U has to be replaced by the *Dirac evolution transport* DU given by

$${}^DU_\gamma(t, s) = l_{\gamma(t)}^{-1} \circ {}^DU(t, s) \circ l_{\gamma(s)}, \quad s, t \in J$$

(cf. (2.4)). The *bundle Dirac equation* is

$${}^DD_t^\gamma \Psi_\gamma = 0 \quad (3.12)$$

with DD being the assigned to DU by (2.7) derivation along paths. Again, the matrix of the coefficients of the Dirac evolution transport is connected with the matrix-bundle Dirac Hamiltonian via (2.10), etc.

3.3. Klein-Gordon equation

The wavefunction $\phi \in K^m$ of spinless special-relativistic particle is a scalar function of class C^m , $m \geq 2$, over the spacetime and satisfies the Klein-Gordon equation [12, chapte 9]. For a particle of mass m and electric charge e in an external electromagnetic field with 4-potential $(\varphi, \mathbf{A})$ it reads [13, chapter XX, § 5, equation (30)]

$$\left[\left(i\hbar \frac{\partial}{\partial t} - e\varphi \right)^2 - c^2 \left(\mathbf{p} - \frac{e}{c} \mathbf{A} \right)^2 \right] \phi = m^2 c^4 \phi. \quad (3.13)$$

This is a second-order linear partial differential equation of type (3.2) with respect to ϕ . Solving it with respect to $\partial^2 \phi / \partial t^2$, we can transform it to equation of type (3.3):

$$\begin{aligned} \frac{\partial^2 \phi}{\partial t^2} &= f_0 \phi + \frac{2e}{i\hbar} \varphi \frac{\partial \phi}{\partial t} \\ f_0 &:= \left[-\frac{c^2}{\hbar^2} \left(\mathbf{p} - \frac{e}{c} \mathbf{A} \right)^2 - \frac{m^2 c^4}{\hbar^2} + \frac{e^2}{\hbar^2} \varphi^2 + \frac{2e}{i\hbar} \varphi \frac{\partial \varphi}{\partial t} \right] \text{id}_{K^m}. \end{aligned} \quad (3.14)$$

As pointed above, there are (infinitely many) different ways to put this equation into Schrödinger-type form. Below we realize three of them, each having applications for different purposes.

The ‘canonical’ way to do this is to define $\psi := (\phi, \partial \phi / \partial t)^\top$ and $K^m := \{\phi: J \rightarrow \mathbb{C}, \phi \text{ is of class } C^m\}$, $m \geq 2$. Then, comparing (3.14) with (3.6), we see that (3.14), and hence (3.13), is equivalent to (3.1) with $\mathcal{H} = {}^{K-G}_c \mathcal{H}$, where the ‘canonical’ Klein-Gordon Hamiltonian is

$${}^{K-G}_c \mathcal{H} := i\hbar \begin{pmatrix} 0 & \text{id}_{K^m} \\ f_0 & \frac{2e}{i\hbar} \varphi \text{id}_{K^{m-1}} \end{pmatrix}. \quad (3.15)$$

It should be note, for a free scalar particle, $(\varphi, \mathbf{A}) = 0$, this is the anti-diagonal matrix operator $\begin{pmatrix} 0 & 1 \\ c^2 \nabla^2 - m^2 c^4 / \hbar^2 & 0 \end{pmatrix} \text{id}_{K^m}$.

Another possibility is to put $\psi = \left(\phi + \frac{i\hbar}{mc^2} \frac{\partial \phi}{\partial t}, \phi - \frac{i\hbar}{mc^2} \frac{\partial \phi}{\partial t} \right)^\top$. This choice is good for investigation of the non-relativistic limit, when $i\hbar \frac{\partial \phi}{\partial t} \approx mc^2 \phi$ [13, chapter XX, § 5], so that in it $\psi \approx (2\phi, 0)^\top$.

Now, as a simple verification proves, the Schrödinger-type form (3.1) of (3.14) is realized for the Hamiltonian

$${}^{\text{K-G}}_{\text{n.r.}}\mathcal{H} := \frac{1}{2} \times \begin{pmatrix} (mc^2 + 2e\varphi)\text{id}_{K^{m-1}} - \hbar^2 f_0/mc^2 & (-mc^2 - 2e\varphi)\text{id}_{K^{m-1}} - \hbar^2 f_0/mc^2 \\ (mc^2 - 2e\varphi)\text{id}_{K^{m-1}} + \hbar^2 f_0/mc^2 & (-mc^2 + 2e\varphi)\text{id}_{K^{m-1}} + \hbar^2 f_0/mc^2 \end{pmatrix}.$$

If the electromagnetic field vanishes, $(\varphi, \mathbf{A}) = 0$, then $f_0 = (c^2 \nabla^2 - m^2 c^4 / \hbar^2) \text{id}_{K^m}$ and

$$\begin{aligned} {}^{\text{K-G}}_{\text{n.r.}}\mathcal{H} &= \frac{1}{2} \left[mc^2 \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} - \frac{\hbar^2}{2m} \nabla^2 \begin{pmatrix} 1 & 1 \\ -1 & -1 \end{pmatrix} \right] \text{id}_{K^{m-1}} \\ &= \left[\left(mc^2 + \frac{\mathbf{p}^2}{2m} \right) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} + \frac{\mathbf{p}^2}{2m} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right] \text{id}_{K^{m-1}} \end{aligned}$$

The third possibility mentioned corresponds to the choice of ψ as a 5×1 matrix:

$$\psi = (mc^2 \phi, \frac{\partial \phi}{\partial t}, \frac{\partial \phi}{\partial x^1}, \frac{\partial \phi}{\partial x^2}, \frac{\partial \phi}{\partial x^3})^\top = (mc^2 \phi, \frac{\partial \phi}{\partial t}, \nabla \phi)^\top.$$

The corresponding Hamiltonian is 5×5 matrix which in the absence of electromagnetic field is [11, chapter 2, sect. 2.1.1] ${}^{\text{K-G}}_5\mathcal{H} = mc^2 \beta + \boldsymbol{\alpha} \cdot \nabla$ (cf. Dirac Hamiltonian (3.10) for $(\varphi, \mathbf{A}) = 0$) where β is 5×5 matrix and $\boldsymbol{\alpha}$ is a 3-vector of 5×5 matrices. The full realization of this procedure and the explicit form of the corresponding 5×5 matrices is given in [14, § 4, sect. 4.4] (see also below Subsect. 4.2, formulae (4.23)).

Choosing some representation of Klein-Gordon equation as first-order (Schrödinger-type) equation, we can in an evident way transfer the bundle formalism to the description of spinless particles.

3.4. Other relativistic wave equations

As we saw in subsections 3.2 and 3.3, the only problem for a bundle reformulation of a wave equation is to rewrite it as a first-order differential equation and to find the corresponding Hamiltonian. Since all relativistic wave equations are of the form of equation (3.2) [12–14], this procedure can successfully be performed for all of them.

For instance, the wavefunction $\psi = (\psi_0, \psi_1, \psi_2, \psi_3)^\top$ of particles with spin 1 is a 4-vector satisfying Klein-Gordon equation⁸ [14, capter I, § 4]. Hence for each component ψ_i , $i = 0, 1, 2, 3$ we can construct the corresponding Hamiltonian ${}^{\text{K-G}}\mathcal{H}_i$ using, e.g., one of the methods described in Subsect. 3.3. Then equation (3.1) holds for a Hamiltonian of the form of a 4×4 diagonal block matrix operator

$${}^{\text{K-G}}\mathcal{H} = \text{diag}({}^{\text{K-G}}\mathcal{H}_0, {}^{\text{K-G}}\mathcal{H}_1, {}^{\text{K-G}}\mathcal{H}_2, {}^{\text{K-G}}\mathcal{H}_3)$$

and the corresponding new wavefunction which is now a 8×1 matrix.

The just said is, of course, valid with respect to electromagnetic field, the 4-potential playing the rôle of wavefunction [14, chapter I, § 5]. It is interesting to be noted that even at a level of classical electrodynamics the Maxwell equations admit Schrödinger-like form. There are different fashions to do this. For a free field, one of them is to put $\psi = (\mathbf{E}, \mathbf{H}, \mathbf{E}, \mathbf{H})^\top$

⁸Here we do not concern the additional conditions like the Lorentz one. They lead to a modification of the equations defining the evolution transport, but this does not change the main ideas. E.g. the most general equation for vectorial mesons is the Proca equation [11, chapter 3, equation (3.132)] $[(\partial^\alpha \partial_\alpha + \frac{m^2 c^4}{\hbar^2}) \eta_{\mu\nu} - \partial_\mu \partial_\nu] \phi^\nu = 0$ on which one usually imposes the additional condition $\partial_\nu \phi^\nu = 0$. (For $m \neq 0$ the last condition is a corollary of the Proca equation; to prove this, simply apply ∂^μ to Proca equation.)

where $\mathbf{E}$ and $\mathbf{H}$ are respectively the electric and magnetic field strengths, and to define the Hamiltonian, e.g., by ${}^{\text{E-M}}\mathcal{H} = i\hbar \begin{pmatrix} 0 & c\text{rot} & 0 & 0 \\ -c\text{rot} & 0 & 0 & 0 \\ 0 & 0 & 0 & \text{div} \\ 0 & 0 & -\text{div} & 0 \end{pmatrix}$.

Almost the same as spin 1 particles is the case of particles with spin 2. Their wavefunction is symmetric tensor field of second rank whose components satisfy also the Klein-Gordon equation [15, p. 24].

The wavefunction of particles with spin 3/2 is a (4-)spin-vector whose ‘vector’ components satisfy the Dirac equation (and some additional conditions) [15, pp. 35–36]. Therefore for them can be applied *mutatis mutandis* the presented in Subsect. 3.2 bundle formalism.

4. Covariant approach

The developed in Sect. 3 bundle formalism for relativistic quantum wave equations has the deficiency that it is not explicitly covariant; so it is not in harmony with the relativistic theory it represents. This is a consequence of the direct applications of the bundle methods developed for the nonrelativistic region, where they work well enough, to the relativistic one. The present section is intended to mend this ‘defect’. Here we develop an appropriate covariant bundle description of relativistic quantum mechanics which corresponds to the natural character of this theory. The difference between the time-dependent and covariant approach is approximately the same as the one between the Hamiltonian and Lagrangian derivation (and forms) of some relativistic wave equation.

4.1. Dirac equation

The covariant Dirac equation [11, sect. 2.1.2], [13, chapter XX, § 8] for a (spin $\frac{1}{2}$) particle with mass m and electric charge e in an external electromagnetic field with 4-potential $\mathcal{A}^\mu$ is

$$(i\hbar\mathcal{D} - mc\mathbb{1}_4)\psi = 0, \quad \mathcal{D} := \gamma^\mu D_\mu, \quad D_\mu := \partial_\mu - \frac{e}{i\hbar c}\mathcal{A}_\mu. \quad (4.1)$$

Here $i \in \mathbb{C}$ is the imaginary unit, $\hbar$ is the Planck constant (divided by 2π), $\mathbb{1}_4 = \text{diag}(1, 1, 1, 1)$ is the 4×4 unit matrix, $\psi := (\psi^0, \psi^1, \psi^2, \psi^3)$ is (the matrix of the components of) a 4-spinor, γ^μ , $\mu = 0, 1, 2, 3$, are the well known Dirac γ -matrices [11–13], and c is the velocity of light in vacuum. Since (4.1) is a first order partial differential equation on the Minkowski 4-dimensional spacetime, it does not admit an evolution operator with respect to the spacetime. More precisely, if $x_1, x_2 \in M$, M being the spacetime, i.e. the Minkowski space M^4 , then there does *not* exist a 4×4 matrix operator (see Appendix B) $\mathcal{U}(x_2, x_1)$ such that

$$\begin{aligned} \psi(x_2) &= \mathcal{U}(x_2, x_1)\psi(x_1). \\ (i\hbar\mathcal{D}_x - mc\mathbb{1}_4)\mathcal{U}(x, x_0) &= 0, \quad \mathcal{U}(x_0, x_0) = \text{id}_{\mathcal{F}}, \quad x, x_0 \in M \end{aligned}$$

where $\mathcal{D} = \mathcal{D} - \frac{e}{i\hbar c}\mathcal{A}$ with $\mathcal{D} := \gamma^\mu \partial / \partial x^\mu$ and $\mathcal{A} := \gamma^\mu \mathcal{A}_\mu$, $\mathcal{F}$ is the space of 4-spinors, and id_X is the identity map of a set X .

By this reason, the methods of previous sections cannot be applied directly to the general 4-dimensional spacetime description of the Dirac equation. An alternative approach to the problem is presented below.

Suppose (F, π, M) is a vector bundle with (total) bundle space F , projection $\pi: F \rightarrow M$, fibre $\mathcal{F}$, and isomorphic fibres $F_x := \pi^{-1}(x)$, $x \in M$. There exist linear isomorphisms $l_x: F_x \rightarrow \mathcal{F}$ which we assume to be diffeomorphisms; so $F_x = l_x^{-1}(\mathcal{F})$ are 4-dimensional vector spaces.

To a state vector (spinor) $\psi(x_0)$ at a fixed point x_0 , we assign a C^1 section⁹ Ψ_{x_0} of (F, π, M) , i.e. $\Psi_{x_0} \in \text{Sec}^1(F, \pi, M)$, by (cf. (2.3))

$$\Psi_{x_0}(x) := l_x^{-1}(\psi(x_0)) \in F_x := \pi^{-1}(x), \quad x \in M. \quad (4.2)$$

Generally $\Psi_{x_0}(x)$ depends on the choice of the point $x_0 \in M$.

Since in (F, π, M) the state of a Dirac particle is described by Ψ_{x_0} , we call it *state section*; resp. (F, π, M) is the *4-spinor bundle*. The description of Dirac particle via Ψ_{x_0} will be called *bundle description*. If it is known, the conventional spinor description is achieved by the spinor

$$\psi(x_0) := l_x(\Psi_{x_0}(x)) \in \mathcal{F}. \quad (4.3)$$

Evidently, we have

$$\Psi_{x_0}(x_2) = L(x_2, x_1)\Psi_{x_0}(x_1), \quad x_2, x_1 \in M. \quad (4.4)$$

with (cf. (2.4))

$$L(y, x) := l_y^{-1} \circ l_x: F_x \rightarrow F_y, \quad x, y \in M. \quad (4.5)$$

Obviously, $\mathcal{L}$ is a linear 4×4 matrix operator satisfying the equations

$$L(x_3, x_1) = L(x_3, x_2) \circ L(x_2, x_1), \quad x_1, x_2, x_3 \in M, \quad (4.6)$$

$$L(x, x) = \text{id}_{F_x}, \quad x \in M. \quad (4.7)$$

Consequently, by definition A.1, the map $L: (y, x) \rightarrow L(y, x) = K_{x \rightarrow y}^{\text{id}_M}$ is a linear transport along the identity map id_M of M in the bundle (F, π, M) . Alternatively, as it is mentioned in Appendix A, this means that $L^\gamma: (t, s) \rightarrow L_{s \rightarrow t}^\gamma := L(\gamma(t), \gamma(s))$, $s, t \in J$ is a flat linear transport along $\gamma: J \rightarrow M$ in (F, π, M) . (Besides, $L(y, x)$ is a Hermitian and unitary transport — see [2].)

Equation (4.4) simply means that Ψ_{x_0} is *L-transported (along id_M) section* of (F, π, M) (cf. [17, definition 5.1]). Writing (A.10) for the transport L and applying the result to a state section given by (4.2), one can prove that (4.4) is equivalent to

$$\mathcal{D}_\mu \Psi = 0, \quad \mu = 0, 1, 2, 3 \quad (4.8)$$

where, for brevity, we have put $\mathcal{D}_\mu := \mathcal{D}_{x^\mu}^{\text{id}_M}$ which is the μ -th partial (section-)derivation along the identity map (of the spacetime) assigned to the transport L .

Now we shall introduce local bases and take a local view of the above-described material.

Let $\{f_\mu(x)\}$ be a basis in $\mathcal{F}$ and $\{e_\mu(x)\}$ be a basis in F_x , $x \in M$. The matrices corresponding to vectors and/or linear maps (operators) in these fields of bases will be denoted by the same (kernel) symbol but in **boldface**, for instance: $\boldsymbol{\psi} := (\psi^0, \psi^1, \psi^2, \psi^3)^\top$ and $\boldsymbol{l}_x(y) := [(l_x)^\mu_\nu(y)]$ are defined, respectively, by $\psi(x) =: \psi^\mu(x)f_\mu(x)$ and $l_x(e_\nu(x)) =: (l_x(y))^\mu_\nu f_\mu(y)$. We put $\boldsymbol{l}_x := \boldsymbol{l}_x(x)$; in fact this will be the only case when the matrix of l_x will be required as we want the ‘physics in F_x ’ to correspond to that of $\mathcal{F}$ at $x \in M$. A very convenient choice is to put $e_\mu(x) = l_x^{-1}(f_\mu(x))$; so then $\boldsymbol{l}_x = \mathbb{1}_4 := [\delta^\mu_\nu] = \text{diag}(1, 1, 1, 1)$ is the 4×4 unit matrix.

The matrix elements of the mapping $L(y, x)$ are defined via the equation $L(y, x)(e_\mu(x)) =: L^\lambda_\mu(y, x)e_\lambda(y)$ and, due to (4.5), we have

$$\boldsymbol{L}(y, x) = \boldsymbol{l}_y^{-1} \cdot \boldsymbol{l}_x \quad (4.9)$$

⁹In contrast to the time-dependent approach (Sect. 3) and nonrelativistic case (Sect. 2) now Ψ_{x_0} is simply a section, not section along paths [2]. Physically this corresponds to the fact that quantum objects do not have world lines (trajectories) in a classical sense [16].

which is generically a matrix operator (see Appendix B).

According to (A.12) the *coefficients* of the transport L form four matrix operators

$${}_{\mu}\mathbf{\Gamma}(x) := [\Gamma^{\lambda}_{\nu}(x)]^3_{\lambda,\nu=0} := \left. \frac{\partial \mathbf{L}(x, y)}{\partial y^{\mu}} \right|_{y=x} = \mathbf{l}^{-1}(x) \frac{\partial \mathbf{l}(x)}{\partial x^{\mu}} \quad (4.10)$$

where (4.9) was applied (see also theorem A.1).

Applying (A.5) and (4.10), we find

$$\frac{\partial \mathbf{L}(y, x)}{\partial y^{\mu}} = -{}_{\mu}\mathbf{\Gamma}(y) \odot \mathbf{L}(y, x), \quad \frac{\partial \mathbf{L}(y, x)}{\partial x^{\mu}} = \mathbf{L}(y, x) \odot {}_{\mu}\mathbf{\Gamma}(x), \quad (4.11)$$

where $\odot$ denotes the introduced by (B.2) multiplication of matrix operators. Therefore

$$\partial_y \mathbf{L}(y, x) = -\mathbf{\Gamma}(y) \mathbf{L}(y, x), \quad \mathbf{\Gamma}(x) := \gamma^{\mu} \cdot {}_{\mu}\mathbf{\Gamma}(x). \quad (4.12)$$

Similarly to the nonrelativistic case [3], to any operator $\mathcal{A}: \mathcal{F} \rightarrow \mathcal{F}$ we assign a bundle morphism $A: F \rightarrow F$ by

$$A_x := A|_{F_x} := l_x^{-1} \circ \mathcal{A} \circ l_x. \quad (4.13)$$

Defining

$$G^{\mu}(x) := l_x^{-1} \circ \gamma^{\mu} \circ l_x, \quad d_{\mu} := d_{\mu}|_x := l_x^{-1} \circ \partial_{\mu} \circ l_x, \quad \partial_{\mu} := \frac{\partial}{\partial x^{\mu}} \quad (4.14)$$

and using the matrices $\mathcal{G}^{\mu}(x)$ and $\mathcal{E}_{\mu}(x)$ given via (B.12), we get

$$\mathbf{G}^{\mu}(x) = l_x^{-1} \mathcal{G}^{\mu}(x) l_x, \quad \mathbf{d}_{\mu} := \mathbb{1}_4 \partial_{\mu} + l_x^{-1} (\partial_{\mu} l_x + \mathcal{E}_{\mu}(x) l_x). \quad (4.15)$$

(Here and below, for the sake of shortness, we sometimes omit the argument x .) The anti-commutation relations

$$\begin{aligned} G^{\mu} G^{\nu} + G^{\nu} G^{\mu} &= 2\eta^{\mu\nu} \text{id}_F, \\ \mathbf{G}^{\mu} \mathbf{G}^{\nu} + \mathbf{G}^{\nu} \mathbf{G}^{\mu} &= \mathcal{G}^{\mu} \mathcal{G}^{\nu} + \mathcal{G}^{\nu} \mathcal{G}^{\mu} = 2\eta^{\mu\nu} \mathbb{1}_4 \quad (= \gamma^{\mu} \gamma^{\nu} + \gamma^{\nu} \gamma^{\mu}), \end{aligned} \quad (4.16)$$

where $[\eta^{\mu\nu}] = \text{diag}(1, -1, -1, -1) = [\eta_{\mu\nu}]$ is the Minkowski metric tensor, can be verified by means of (4.14), (4.15), (B.10), and the well know analogous relation for the γ -matrices (see, e.g. [12, chapter 2, equation (2.5)]).

For brevity, if $a_{\mu}: F \rightarrow F$ are morphisms, sums like $G^{\mu} \circ a_{\mu}$ will be denoted by ‘back-slashing’ the kernel letter, $\mathfrak{d} := G^{\mu} \circ a_{\mu}$ (cf. the ‘slashed’ notation $\not{a} := \gamma^{\mu} a_{\mu}$). Similarly, we put $\mathfrak{d} := \mathbf{G}^{\mu}(x) a_{\mu}$ for $a_{\mu} \in GL(4, \mathbb{C})$. It is almost evident (see (4.14)) that the morphism corresponding to $\not{a} := \gamma^{\mu} a_{\mu}$ is $\mathfrak{d} = G^{\mu}(x) \circ d_{\mu}$:

$$\mathfrak{d} = l_x^{-1} \circ \not{a} \circ l_x. \quad (4.17)$$

Now we can write the Dirac equation (4.1) in a bundle form.

First of all, we rewrite (4.1) as

$$i\hbar \not{\partial}_x \psi(x) = \mathcal{D}_x \psi(x), \quad \mathcal{D}_x := mc \mathbb{1}_4 + \frac{e}{c} \mathcal{A}(x), \quad (4.18)$$

the index x meaning that the corresponding operators act with respect to the variable $x \in M$. This is the *covariant* Schrödinger-like form of Dirac equation; $\not{\partial}_x$ is the analogue of the time derivation d/dt and $\mathcal{D}_x$ corresponds to the Hamiltonian $\mathcal{H}$. We call $\mathcal{D}$ the *Dirac function*, or simply, *Diracian* of a particle described by Dirac equation.

Substituting (4.3) into (4.18), acting on the result from the left by l_y^{-1} , and using (4.14), we find the bundle form of (4.18) as

$$i\hbar \mathbb{d}_x|_y \Psi_x(y) = D_x|_y \Psi_x(y) \quad (4.19)$$

with $y \in M$, $\mathbb{d}_x|_y := G(y)d_\mu|_x|_y$, $d_\mu|_x|_y := l_y^{-1} \circ \frac{\partial}{\partial x^\mu} \circ l_y$, $D_x \in \text{Mor}(F, \pi, M)$ being the (Dirac) bundle morphism assigned to the Diracian. We call it the *bundle Diracian*. According to (4.13) it is defined by

$$D_x|_y := D_x|_{F_y} = l_y^{-1} \circ \mathcal{D}_x \circ l_y = mc \text{id}_{F_y} + \frac{e}{c} \mathbb{A}_x|_y, \quad (4.20)$$

where $A_\mu|_x = \mathcal{A}_\mu|_x \text{id}_F \in \text{Mor}(F, \pi, M)$, $A_\mu|_x|_y = l_y^{-1} \circ (\mathcal{A}_\mu|_x \text{id}_F) \circ l_y = \mathcal{A}_\mu|_x \text{id}_{F_y}$ are the components of the *bundle electromagnetic potential* and

$$\mathbb{A}_x|_y = G^\mu(y) \circ A_\mu|_x|_y = G^\mu(y) \circ (l_y^{-1} \circ \mathcal{A}_\mu \text{id}_F \circ l_y) = l_y^{-1} \circ (\mathcal{A}_\mu|_x \text{id}_F) \circ l_y = \mathcal{A}_x|_y. \quad (4.21)$$

4.2. Other relativistic wave equations

The relativistic-covariant Klein-Gordon equation for a (spinless) particle of mass m and electric charge e in a presence of (external) electromagnetic field with 4-potential $\mathcal{A}_\mu$ is [13, chapter XX, § 5, equation (30')]

$$\left(\mathcal{D}^\mu \mathcal{D}_\mu + \frac{m^2 c^2}{\hbar^2} \right) \phi = 0, \quad \mathcal{D}_\mu = \eta_{\mu\nu} \mathcal{D}^\nu := \partial_\mu - \frac{e}{i\hbar c} \mathcal{A}_\mu. \quad (4.22)$$

Since this is a second-order partial differential equation, it does not directly admit an evolution operator and adequate bundle formulation and interpretation. To obtain such a formulation, we have to rewrite (4.22) as a first-order (system of) partial differential equation(s) (cf. Subsect. 3.3).

Perhaps the best way to do this is to replace ϕ with a 5×1 matrix $\varphi = (\varphi^0, \dots, \varphi^4)^\top$ and to introduce 5×5 Γ^μ , $\mu = 0, 1, 2, 3$ with components $(\Gamma^\mu)^i_j$, $i, j = 0, 1, 2, 3, 4$ such that (cf. [14, chapter I, equations (4.38) and (4.37)]):

$$\varphi = a \begin{pmatrix} i\hbar \mathcal{D}_0 \phi \\ i\hbar \mathcal{D}_1 \phi \\ i\hbar \mathcal{D}_2 \phi \\ i\hbar \mathcal{D}_3 \phi \\ mc \phi \end{pmatrix}, \quad (\Gamma^\mu)^i_j = \begin{cases} 1 & \text{for } (i, j) = (\mu, 4) \\ \eta_{\mu\mu} & \text{for } (i, j) = (4, \mu) \\ 0 & \text{otherwise} \end{cases} \quad (4.23)$$

where the complex constant $a \neq 0$ is insignificant for us and can be (partially) fixed by an appropriate normalization of φ .

Then a simple checking shows that (4.22) is equivalent to (cf. (4.1))

$$(i\hbar \Gamma^\mu \mathcal{D}_\mu - mc \mathbb{1}_5) \varphi = 0 \quad (4.24)$$

with $\mathbb{1}_5 = \text{diag}(1, 1, 1, 1, 1)$ being the 5×5 unit matrix, or (cf. (4.18))

$$i\hbar \Gamma^\mu \partial_\mu \varphi = \mathcal{K} \varphi, \quad \mathcal{K} := mc \mathbb{1}_5 + \frac{e}{c} \Gamma^\mu \mathcal{A}_\mu. \quad (4.25)$$

Now it is evident that *mutatis mutandis*, taking Γ^μ for γ^μ , φ for ψ , etc., the (bundle) machinery developed in Subsect. 4.1 for the Dirac equation can be applied to the Klein-Gordon equation in the form (4.24). Since the transferring of the results obtained in Subsect. 4.1 for

Dirac equation to Klein-Gordon one is absolutely trivial,¹⁰ we are not going to present here the bundle description of the latter equation.

Since the relativistic wave equations for particles with spin greater than 1/2 are versions or combinations of Dirac and Klein-Gordon equations [12–15], for them is *mutatis mutandis* applicable the bundle approach developed in Subsect. 4.1 for Dirac equation or/and its version for Klein-Gordon one pointed above.

5. Propagators and evolution transports or operators

The propagators, called also propagator functions or Green functions, are solutions of the wave equations with point-like unit source and satisfy appropriate (homogeneous) boundary conditions corresponding to a concrete problem under exploration [11, 12]. Undoubtedly these functions play an important rôle in the mathematical apparatus of (relativistic) quantum mechanics and its physical interpretation [11, 12]. By this reason it is essential to be investigated the connection between propagators and evolution operators or/and transports. As we shall see below, the latter can be represented as integral operators whose kernel is connected in a simple way with the corresponding propagator.

5.1. Green functions (review)

Generally [18, article “Green function”] the *Green function* $g(x', x)$ of a linear differential operator L (or of the equation $Lu(x) = f(x)$) is the kernel of the integral operator inverse to L . As the kernel of the unit operator is the Dirac delta-function $\delta^4(x' - x)$, the Green function is a fundamental solution of the non-homogeneous equation

$$Lu(x) = f(x), \quad (5.1)$$

i.e. treated as a generalized function $g(x', x)$ is a solution of

$$L_{x'}g(x', x) = \delta^4(x' - x). \quad (5.2)$$

Given a Green function $g(x', x)$, the solution of (5.1) is

$$u(x') = \int g(x', x)f(x) d^4x. \quad (5.3)$$

A concrete Green function $g(x', x)$ for L (or (5.1)) satisfies, besides (5.2), certain (homogeneous) boundary conditions on x' with fixed x , i.e. it is the solution of a fixed boundary-value problem for equation (5.2). Hence, if $g_f(x', x)$ is some fundamental solution, then

$$g(x', x) = g_f(x', x) + g_0(x', x) \quad (5.4)$$

where $g_0(x', x)$ is a solution of the homogeneous equation $L_{x'}g_0(x', x) = 0$ chosen such that $g(x', x)$ satisfies the required boundary conditions.

Suppose $g(x', x)$ is a Green function of L for some boundary-value (or initial-value) problem. Then, using (5.2), we can verify that

$$L_{x'} \left(\int g(x', x)u(x) d^3x \right) = \delta(ct' - ct)u(x') \quad (5.5)$$

¹⁰Both coincide up to notation or a meaning of the corresponding symbols.

where $x = (ct, \mathbf{x})$ and $x' = (ct', \mathbf{x}')$. Therefore the solution of the problem

$$L_x u(x) = 0, \quad u(ct_0, \mathbf{x}) = u_0(\mathbf{x}) \quad (5.6)$$

is

$$u(x) = \int g(x, (ct_0, \mathbf{x}_0)) u_0(\mathbf{x}_0) d^3 \mathbf{x}_0, \quad x_0 = (ct_0, \mathbf{x}_0) \quad \text{for } t \neq t_0. \quad (5.7)$$

From here we can make the conclusion that, if (5.6) admits an evolution operator $\mathcal{U}$ such that (cf. (2.2))

$$u(x) \equiv u(ct, \mathbf{x}) = \mathcal{U}(t, t_0) u(ct_0, \mathbf{x}), \quad (5.8)$$

then the r.h.s of (5.7) realizes $\mathcal{U}$ as an integral operator with a kernel equal to the Green function g .

Since all (relativistic or not) wave equations are versions of (5.6), the corresponding evolution operators, if any, and Green functions (propagators) are connected as just described. Moreover, if some wave equation does not admit (directly) evolution operator, e.g. if it is of order greater than one, then we can *define* it as the corresponding version of the integral operator in the r.h.s. of (5.7). In this way is established a one-to-one onto correspondence between the evolution operators and Green functions for any particular problem like (5.6).

And a last general remark. The so-called S-matrix finds a lot of applications in quantum theory [11, 12, 14]. By definition S is an operator transforming the system's state vector $\psi(-\infty, \mathbf{x})$ before scattering (reaction) into the one $\psi(+\infty, \mathbf{x})$ after it:

$$\lim_{t \rightarrow +\infty} \psi(ct, \mathbf{x}) =: S \lim_{t \rightarrow -\infty} \psi(ct, \mathbf{x}).$$

So, e.g. when (5.8) takes place, we have

$$S = \lim_{t_{\pm} \rightarrow \pm\infty} \mathcal{U}(t_+, t_-) =: \mathcal{U}(+\infty, -\infty). \quad (5.9)$$

Thus the above-mentioned connection between evolution operators and Green functions can be used for expressing the S-matrix in terms of propagators. Such kind of formulae are often used in relativistic quantum mechanics [12].

5.2. Nonrelativistic case (Schrödinger equation)

The (retarded) Green function $g(x', x)$, $x', x \in M$, for the Schrödinger equation (2.1) is defined as the solution of the boundary-value problem [12, § 22]

$$\left[i\hbar \frac{\partial}{\partial t'} - \mathcal{H}(x') \right] g(x', x) = \delta^4(x' - x), \quad (5.10)$$

$$g(x', x) = 0 \quad \text{for } t' < t \quad (5.11)$$

where $x' = (ct', \mathbf{x}')$, $x = (ct, \mathbf{x})$, $\mathcal{H}(x)$ is system's Hamiltonian, and $\delta^4(x' - x)$ is the 4-dimensional (Dirac) δ -function.

Given g , the solution $\psi(x')$ (for $t' > t$) of (2.1) is¹¹

$$\theta(t' - t) \psi(x') = i\hbar \int d^3 \mathbf{x} g(x', x) \psi(x) \quad (5.12)$$

¹¹One should not confuse the notation $\psi(x) = \psi(ct, \mathbf{x})$, $x = (ct, \mathbf{x})$ of this section and $\psi(t)$ from Sect. 2. The latter is the wavefunction at a moment t and the former is its value at the spacetime point $x = (ct, \mathbf{x})$. Analogously, $\Psi_\gamma(x) \equiv \Psi_\gamma(ct, \mathbf{x}) := l_{\gamma(t)}^{-1}(\psi(ct, \mathbf{x}))$ should not be confused with $\Psi_\gamma(t)$ from Sect. 2. A notation like $\psi(t)$ and $\Psi_\gamma(t)$ will be used if the spatial parts of the arguments are inessential, as in Sect. 2, and there is no risk of ambiguities.

where the θ -function $\theta(s)$, $s \in \mathbb{R}$, is defined by $\theta(s) = 1$ for $s > 0$ and $\theta(s) = 0$ for $s < 0$.

Combining (2.2) and (5.12), we find the basic connection between the evolution operator $\mathcal{U}$ and Green function of Schrödinger equation:

$$\theta(t' - t)[\mathcal{U}(t', t)(\psi(ct, \mathbf{x}'))] = i\hbar \int d^3\mathbf{x} g((ct', \mathbf{x}'), (ct, \mathbf{x}))\psi(ct, \mathbf{x}). \quad (5.13)$$

Actually this formula, if g is known, determines $\mathcal{U}(t', t)$ for all t' and t , not only for $t' > t$, as $\mathcal{U}(t', t) = \mathcal{U}^{-1}(t, t')$ and $\mathcal{U}(t, t) = \text{id}_{\mathcal{F}}$ with $\mathcal{F}$ being the system's Hilbert space (see (2.2) or [2, sect. 2]). Consequently *the evolution operator for the Schrödinger equation can be represented as an integral operator whose kernel, up to the constant $i\hbar$, is exactly the (retarded) Green function for it.*

To write the bundle version of (5.13), we introduce the *Green operator* which is simply a multiplication with the Green function:

$$\mathcal{G}(x', x) := g(x', x) \text{id}_{\mathcal{F}}: \mathcal{F} \rightarrow \mathcal{F}. \quad (5.14)$$

The corresponding to it *Green morphism* G is given via (see (2.11) and cf. (2.4))

$$G_{\gamma}(x', x) := l_{\gamma(t')}^{-1} \circ \mathcal{G}(x', x) \circ l_{\gamma(t)} = g(x', x) l_{\gamma(t')}^{-1} \circ l_{\gamma(t)}: F_{\gamma(t)} \rightarrow F_{\gamma(t')}. \quad (5.15)$$

Now, acting on (5.13) from the left by $l_{\gamma(t')}^{-1}$ and using (2.3) and (2.4), we obtain

$$\theta(t' - t)[U_{\gamma}(t', t)(\Psi_{\gamma}(ct, \mathbf{x}'))] = i\hbar \int d^3\mathbf{x} G_{\gamma}((ct', \mathbf{x}'), (ct, \mathbf{x}))\Psi_{\gamma}(ct, \mathbf{x}). \quad (5.16)$$

Therefore for the Schrödinger equation the *evolution transport* U can be represented as an *integral operator with kernel equal to $i\hbar$ times the Green morphism G .*

Taking as a starting point (5.13) and (5.16), we can obtain different representations for the evolution operator and transport by applying concrete formulae for the Green function. For example, if a complete set $\{\psi_a(x)\}$ of orthonormal solutions of Schrödinger equation satisfying the completeness condition¹²

$$\sum_a \psi_a(ct, \mathbf{x}') \psi_a^*(ct, \mathbf{x}) = \delta^3(\mathbf{x}' - \mathbf{x})$$

is known, then [12, § 22]

$$g(x', x) = \frac{1}{i\hbar} \theta(t' - t) \sum_a \psi_a(x') \psi_a^*(x)$$

which, when substituted into (5.13), implies

$$\mathcal{U}(t', t)\psi(ct, \mathbf{x}') = \sum_a \psi_a(x') \int d^3\mathbf{x} \psi_a^*(ct, \mathbf{x})\psi(ct, \mathbf{x}). \quad (5.17)$$

Note, the integral in this equation is equal to the a -th coefficient of the expansion of ψ over $\{\psi_a\}$.

¹²Here and below the symbol $\sum_a$ denotes a sum and/or integral over the discrete and/or continuous spectrum. The asterisk (*) means complex conjugation.

5.3. Dirac equation

Since the Dirac equation (4.1) is a first-order linear partial differential equation, it admits both evolution operator and Green function(s) (propagator(s)). From a generic view-point, the only difference from the Schrödinger equation is that (4.1) is a *matrix* equation; so the corresponding Green functions are actually Green matrices, i.e. Green matrix-valued functions. Otherwise the results of Subsect. 5.2 are *mutatis mutandis* applicable to the theory of Dirac equation.

The (retarded) Green matrix (function) or propagator for Dirac equation (4.1) is a 4×4 matrix-valued function $g(x', x)$ depending on two arguments $x', x \in M$ and such that [11, sect. 2.5.1 and 2.5.2]

$$(i\hbar\mathcal{D}_{x'} - mc\mathbf{1}_4)g(x', x) = \delta(x' - x) \quad (5.18)$$

$$g(x', x) = 0 \quad \text{for } t < t'. \quad (5.19)$$

For a free Dirac particle, i.e. for $\mathcal{D} = \partial$ or $\mathcal{A}_\mu = 0$, the explicit expression $g_0(x', x)$ for $g(x', x)$ is derived in [11, sect. 2.5.1], where the notation $\mathcal{K}$ instead of g_0 is used. In an external electromagnetic field $\mathcal{A}_\mu$ the Green matrix g is a solution of the integral equation¹³

$$g(x', x) = g_0(x', x) + \int d^4y g_0(x', y) \frac{e}{c} \mathcal{A}(y) g(y, x) \quad (5.20)$$

which includes the corresponding boundary condition.¹⁴ The iteration of this equation results in the perturbation series for g (cf. [11, sect. 2.5.2]).

If the (retarded) Green matrix $g(x', x)$ is known, the solution $\psi(x')$ of Dirac equation (for $t' > t$) is

$$\theta(t' - t)\psi(x') = i\hbar \int d^3\mathbf{x} g(x', x) \gamma^0 \psi(x). \quad (5.21)$$

Hence, denoting by $\mathcal{U}$ the non-relativistic (see Subsect. 3.2) Dirac evolution operator, from the equations (2.2), and (5.21), we find:

$$\theta(t' - t)[\mathcal{U}(t', t)(\psi(ct, \mathbf{x}'))] = i\hbar \int d^3\mathbf{x} g((ct', \mathbf{x}'), (ct, \mathbf{x})) \gamma^0 \psi(ct, \mathbf{x}). \quad (5.22)$$

So, the evolution operator admits an integral representation whose kernel, up to the right multiplication with $i\hbar\gamma^0$, is equal to the (retarded) Green function for Dirac equation.

Similarly to (5.16), now the bundle version of (5.22) is

$$\theta(t' - t)[U(t', t)(\Psi_\gamma(ct, \mathbf{x}'))] = i\hbar \int d^3\mathbf{x} G_\gamma(x', x) G^0(\gamma(t)) \Psi_\gamma(ct, \mathbf{x}), \quad (5.23)$$

where $G^0(x)$ is defined by (4.14) with $\mu = 0$ and (cf. (5.15))

$$G_\gamma(x', x) := l_{\gamma(t')}^{-1} \circ \mathcal{G}(x', x) \circ l_{\gamma(t)}, \quad G(x', x) := l_{x'}^{-1} \circ \mathcal{G}(x', x) \circ l_x \quad (5.24)$$

with (cf. (5.14))

$$\mathcal{G}(x', x) = g(x', x) \text{id}_{\mathcal{F}}. \quad (5.25)$$

(Here $\mathcal{F}$ is the space of 4-spinors.)

Analogously to the above results, one can obtain such for other propagators, e.g. for the Feynman one [11, 12], but we are not going to do this here as it is a trivial variant of the procedure described.

¹³The derivation of (5.20) is the same as for the Feynman propagators S_F and S_A given in [11, sect. 2.5.2]. The propagators S_F and S_A correspond to g_0 and g respectively, but satisfy other boundary conditions [11, 12].

¹⁴Due to (5.4), the integral equation (5.20) is valid for any Green function (matrix) of the Dirac equation (4.1).

5.4. Klein-Gordon equation

The (retarded) Green function $g(x', x)$ for the Klein-Gordon equation (4.22) is a solution to the boundary-value problem [11, sect. 1.3.1]

$$\left(\mathcal{D}^\mu \mathcal{D}_\mu + \frac{m^2 c^2}{\hbar^2}\right)\Big|_{x'} g(x', x) = \delta^4(x' - x), \quad (5.26)$$

$$g(x', x) = 0 \quad \text{for } t' < t. \quad (5.27)$$

For a free particle its explicit form can be found in [11, sect. 1.3.1].

A simple verification proves that, if $g(x', x)$ is known, the solution ϕ of (4.22) (for $t' > t$) is given by ($x^0 = ct$)

$$\theta(t' - t)\phi(x') = \int d^3\mathbf{x} \left[\frac{\partial g(x', x)}{\partial x^0} \phi(x) + g(x', x) \left(2 \frac{\partial \phi(x)}{\partial x^0} - \frac{e}{i\hbar c} \mathcal{A}^0(x) \phi(x) \right) \right]. \quad (5.28)$$

Introducing the matrices

$$\psi(x) := \begin{pmatrix} \phi(x) \\ \partial_0|_x \phi(x) \end{pmatrix}, \quad \mathbf{g}(x', x) := \begin{pmatrix} \mathcal{D}_0|_x g(x', x) \\ 2g(x', x) \end{pmatrix}, \quad (5.29)$$

we can rewrite (5.28) as

$$\theta(t' - t)\phi(x') = \int d^3\mathbf{x} \mathbf{g}^\top(x', x) \cdot \psi(x) \quad (5.30)$$

where the dot $(\cdot)$ denotes matrix multiplication.

An important observation is that for ψ the Klein-Gordon equation transforms into first-order Schrödinger-type equation (see Subsect. 3.3) with Hamiltonian ${}^{\text{K-G}}\mathcal{H}$ given by (3.15) in which id_{ldots} is replaced by $c\text{id}_{ldots}$.

Denoting the (retarded) Green function, which is in fact 2×2 matrix, and the evolution operator for this equation by $\tilde{\mathcal{G}}(x', x) = [\tilde{\mathcal{G}}_b^a(x', x)]_{a,b=1}^2$ and $\tilde{\mathcal{U}}(t', t) = [\tilde{\mathcal{U}}_b^a(t', t)]_{a,b=1}^2$, respectively, we see that (cf. Subsect. 5.2, equation (5.12))

$$\theta(t' - t)[\tilde{\mathcal{U}}(t', t)\psi(ct, \mathbf{x})] = \theta(t' - t)\psi(x') = \int d^3\mathbf{x} \tilde{\mathcal{G}}(x', x)\psi(x). \quad (5.31)$$

Comparing this equations with (5.29) and (5.30), we find

$$(\tilde{\mathcal{G}}_1^1(x', x), \tilde{\mathcal{G}}_2^1(x', x)) = \mathbf{g}^\top(x', x).$$

The other matrix elements of $\tilde{\mathcal{G}}$ can also be connected with $\mathbf{g}(x', x)$ and its derivatives, but this is inessential for the following.

In this way we have connected, via (5.31), the evolution operator and the (retarded) Green function for a concrete first-order realization of Klein-Gordon equation. It is almost evident that this procedure *mutatis mutandis* works for any such realization; in every case the corresponding Green function (resp. matrix) being a (resp. matrix-valued) function of the Green function $g(x', x)$ introduce via (5.26) and (5.27). For instance, the treatment of the 5-dimensional realization given by (4.23) and (4.24) is practically identical to the one of Dirac equation in Subsect. 5.3, only the γ -matrices γ^μ have to be replace with the 5×5 matrices Γ^μ (defined by (4.23)). This results in a 5×5 matrix evolution operator $\mathcal{U}(x', x)$, etc.

Since the bundle version of (5.31) or an analogous result for $\mathcal{U}(x', x)$ is absolutely trivial (cf. Subsect. 5.2 and 5.3 resp.), we are not going to write it here; up to the meaning of notation it coincides with (5.16) or (5.23) respectively.

6. Conclusion

In this investigation we have reformulated the relativistic wave equations in terms of fibre bundles. In the bundle formulation the wavefunctions are represented as (state) liftings of paths (equivalently: sections along paths) (time-dependent approach) or simply sections (covariant approach) of a suitable vector bundle over the spacetime. The covariant approach has an advantage of being explicitly covariant while in the time-dependent one the time plays a privilege rôle. In both cases the evolution (in time or in spacetime resp.) is described via a linear transport in the bundle mentioned. The state liftings/sections are linearly transported by means of the corresponding (evolution) transports. We also have explored some links between evolution operators or transports and the retarded Green functions (or matrices) for the corresponding wave equations: the former turn to have realization as integral operators whose kernel is equal to the latter ones up to a multiplication with a constant complex number or matrix.

These connections suggest the idea for introducing ‘retarded’, or, in a sense, ‘causal’ evolution operators or transports as a product of the evolution operators or transports with θ -function of the difference of the times corresponding to the first and second arguments of the transport or operator.

Most of the possible generalizations of the bundle non-relativistic quantum mechanics, pointed in [6], are *mutatis mutandis* valid with respect to the bundle version of relativistic quantum mechanics, developed in the present work. The only essential change is that, in the relativistic region, the spacetime model is fixed as the Minkowski spacetime.

A further development of the ideas presented in this paper leads to their application to (quantum) field theory which will be done elsewhere.

We shall end with some comments on the material of Subsect. 3.1. As we saw there, the fibre bundle formalism developed for the solutions of Schrödinger equation can successfully be applied for the solutions of (systems of) linear ordinary differential equations. For this purpose the system of equations, if they are of order greater than one, has to be transformed into a system of first-order equations. It can always be written in a Schrödinger-like form (3.1) to which the developed in [2–6] bundle approach can be applied *mutatis mutandis*.

Therefore, in particular, to a system of linear ordinary (with respect to ‘time’) differential equations corresponds a suitable linear transport along paths in an appropriately chosen fibre bundle. As most of the fundamental equations of physics are expressed by such systems of equations, they admit fibre bundle (re)formulation analogous to the one of Schrödinger equation.

An interesting consequence of this discussion is worth mentioning. Suppose a system of the above-described type is an Euler-Lagrange (system of) equation(s) for some Lagrangian. Applying the outlined ‘bundle’ procedure, we see that to this Lagrangian corresponds some (evolution) linear transport along paths in a suitable fibre bundle. The bundle and the transport are practically (up to isomorphisms) unique if the Euler-Lagrange equations are of first order with respect to time. Otherwise there are different (but equivalent) such objects corresponding to the given Lagrangian. Hence, some Lagrangians admit description in terms of linear transports along paths. In more details the correspondence between Lagrangians (or Hamiltonians) and linear transports along paths will be explored elsewhere.

Appendix A. Linear transports along maps in fibre bundles

In this appendix we recall a few simple facts concerning (linear) transports along maps, in particular along paths, required for the present investigation. The below-presented material

is abstracted from [9, 10, 19] where further details can be found (see also [2, sect. 3]).

Let (E, π, B) be a topological bundle with base B , bundle (total, fibre) space E , projection $\pi : E \rightarrow B$, and homeomorphic fibres $\pi^{-1}(x)$, $x \in B$. Let the set N be not empty, $N \neq \emptyset$, and there be given a map $\varkappa : N \rightarrow B$. By id_X is denoted the identity map of a set X .

Definition A.1. A transport along maps in the bundle (E, π, B) is a map K assigning to any map $\varkappa : N \rightarrow B$ a map $K^\varkappa$, transport along $\varkappa$, such that $K^\varkappa : (l, m) \mapsto K_{l \rightarrow m}^\varkappa$, where for every $l, m \in N$ the map

$$K_{l \rightarrow m}^\varkappa : \pi^{-1}(\varkappa(l)) \rightarrow \pi^{-1}(\varkappa(m)), \quad (\text{A.1})$$

called transport along $\varkappa$ from l to m , satisfies the equalities:

$$K_{m \rightarrow n}^\varkappa \circ K_{l \rightarrow m}^\varkappa = K_{l \rightarrow n}^\varkappa, \quad l, m, n \in N, \quad (\text{A.2})$$

$$K_{l \rightarrow l}^\varkappa = \text{id}_{\pi^{-1}(\varkappa(l))}, \quad l \in N. \quad (\text{A.3})$$

If (E, π, B) is a complex (or real) vector bundle and the maps (A.1) are linear, i.e.

$$K_{l \rightarrow m}^\varkappa(\lambda u + \mu v) = \lambda K_{l \rightarrow m}^\varkappa u + \mu K_{l \rightarrow m}^\varkappa v, \quad \lambda, \mu \in \mathbb{C} \text{ (or } \mathbb{R}), \quad u, v \in \pi^{-1}(\varkappa(l)), \quad (\text{A.4})$$

the transport K is called *linear*. If $\varkappa$ belongs to the set of paths in B , $\varkappa \in \{\gamma : J \rightarrow B, J \text{ being } \mathbb{R}\text{-interval}\}$, we said that the transport K is *along paths*.

For the present work is important that the class of linear transports along the identity map id_B of B coincides with the class of *flat* linear transports along paths¹⁵ (see the comments after equation (2.3) of [19]).

The general form of a transport along maps is described by the following result.

Theorem A.1. Let $\varkappa : N \rightarrow B$. The map $K : \varkappa \mapsto K^\varkappa : (l, m) \mapsto K_{l \rightarrow m}^\varkappa$, $l, m \in N$ is a transport along $\varkappa$ if and only if there exist a set Q and a family of bijective maps $\{F_n^\varkappa : \pi^{-1}(\varkappa(n)) \rightarrow Q, n \in N\}$ such that

$$K_{l \rightarrow m}^\varkappa = (F_m^\varkappa)^{-1} \circ (F_l^\varkappa), \quad l, m \in N. \quad (\text{A.5})$$

The maps $F_n^\varkappa$ are defined up to a left composition with bijective map depending only on $\varkappa$, i.e. (A.5) holds for given families of maps $\{F_n^\varkappa : \pi^{-1}(\varkappa(n)) \rightarrow Q, n \in N\}$ and $\{F'_n : \pi^{-1}(\varkappa(n)) \rightarrow Q', n \in N\}$ for some sets Q and Q' iff there is a bijective map $D^\varkappa : Q \rightarrow Q'$ such that

$$F'_n = D^\varkappa \circ F_n^\varkappa, \quad n \in N. \quad (\text{A.6})$$

For the purposes of this investigation we need a slight generalization of [19, definition 4.1], viz. we want to replace in it $\mathbf{N} \subset \mathbb{R}^k$ with an arbitrary differentiable manifold. Let N be a differentiable manifold and $\{x^a : a = 1, \dots, \dim N\}$ be coordinate system in a neighborhood of $l \in N$. For $\varepsilon \in (-\delta, \delta) \subset \mathbb{R}$, $\delta \in \mathbb{R}_+$ and $l \in N$ with coordinates $l^a = x^a(l)$, we define $l_b(\varepsilon) \in N$, $b = 1, \dots, \dim N$ by $l_b^a(\varepsilon) := x^a(l_b(\varepsilon)) := l^a + \varepsilon \delta_b^a$ where the Kroneker δ -symbol is given by $\delta_b^a = 0$ for $a \neq b$ and $\delta_b^a = 1$ for $a = b$. Let $\xi = (E, \pi, B)$ be a vector bundle, $\varkappa : N \rightarrow B$ be injective (i.e. 1:1 mapping), and $\text{Sec}^p(\xi)$ (resp. $\text{Sec}(\xi)$) be the set of C^p (resp. all) sections over ξ . Let $L_{l \rightarrow m}^\varkappa$ be a C^1 (on l) linear transport along $\varkappa$. Now the modified definition reads:¹⁶

¹⁵The *flat* linear transports along paths are defined as ones with vanishing curvature operator [20, sect. 2]. By [20, theorem. 6.1] we can equivalently define them by the property that they depend only on their initial and final points, i.e. if $K_{s \rightarrow t}^\gamma$, $\gamma : J \rightarrow B$, $s, t \in J \subset \mathbb{R}$, depends only on $\gamma(s)$ and $\gamma(t)$ but not on the path γ itself.

¹⁶We present below, in definition A.2, directly the definition of a *section*-derivation along injective mapping $\varkappa$ as only it will be employed in the present paper (for $\varkappa = \text{id}_N$). If $\varkappa$ is not injective, the mapping (A.7) could be multiple-valued at the points of self-intersection of $\varkappa$, if any. For some details when $\varkappa$ is an arbitrary path in N , see [2, subsect. 3.3].

Definition A.2. The a -th, $1 \leq a \leq \dim N$, partial (section-)derivation along maps generated by L is a map ${}_a\mathcal{D} : \varkappa \mapsto {}_a\mathcal{D}^\varkappa$ where the a -th (partial) derivation ${}_a\mathcal{D}^\varkappa$ along $\varkappa$ (generated by L) is a map

$${}_a\mathcal{D}^\varkappa : l \mapsto \mathcal{D}_{l^a}^\varkappa, \quad (\text{A.7})$$

where the a -th (partial) derivative $\mathcal{D}_{l^a}^\varkappa$ along $\varkappa$ at l is a map

$$\mathcal{D}_{l^a}^\varkappa : \text{Sec}^1 \left(\xi|_{\varkappa(N)} \right) \rightarrow \pi^{-1}(\varkappa(l)) \quad (\text{A.8})$$

defined for $\sigma \in \text{Sec}^1 \left(\xi|_{\varkappa(N)} \right)$ by

$$({}_a\mathcal{D}^\varkappa \sigma)(\varkappa(l)) := \lim_{\varepsilon \rightarrow 0} \left[\frac{1}{\varepsilon} (L_{l^a(\varepsilon) \rightarrow l}^\varkappa \sigma(\varkappa(l_a(\varepsilon))) - \sigma(\varkappa(l))) \right]. \quad (\text{A.9})$$

Accordingly can be modified the other definitions of [19, sect. 4], all the results of it being *mutatis mutandis* valid. In particular, we have:

Proposition A.1. *The operators $\mathcal{D}_{l^a}^\varkappa$ are $(\mathbb{C}-)$ linear and*

$$\mathcal{D}_{m^a}^\varkappa \circ L_{l \rightarrow m}^\varkappa \equiv 0. \quad (\text{A.10})$$

Proposition A.2. *If $\sigma \in \text{Sec}^1 \left(\xi|_{\varkappa(N)} \right)$, then*

$$\mathcal{D}_{l^a}^\varkappa \sigma = \sum_i \left[\frac{\partial \sigma^i(\varkappa(l))}{\partial l^a} + \sum_j {}_a\Gamma_j^i(l; \varkappa) \sigma^j(\varkappa(l)) \right] e_i(l), \quad (\text{A.11})$$

where $\{e_i(l)\}$ is a basis in $\pi^{-1}(\varkappa(l))$, $\sigma(\varkappa(l)) =: \sum_i \sigma^i(\varkappa(l)) e_i(l)$, and the **coefficients** of L are defined by

$${}_a\Gamma_j^i(l; \varkappa) := \left. \frac{\partial L_j^i(l, m; \varkappa)}{\partial m^a} \right|_{m=l} = - \left. \frac{\partial L_j^i(m, l; \varkappa)}{\partial m^a} \right|_{m=l}. \quad (\text{A.12})$$

Here $L_j^i(\cdots)$ are the components of L , $L_{l \rightarrow m}^\varkappa e_i(l) =: \sum_j L_j^i(m, l; \varkappa) e_j(m)$.

The above general definitions and results will be used in this work in the special cases of linear transports along paths and linear transports along the identity map of the bundle's base.

Appendix B. Matrix operators

In this appendix we point to some peculiarities of linear (matrix) operators acting on $n \times 1$, $n \in \mathbb{N}$, matrix fields over the space-time M . Such operators appear naturally in the theory of Dirac equation where one often meets 4×4 matrices whose elements are operators; e.g. an operator of this kind is $\not{\partial} := \gamma^\mu \partial_\mu = [(\gamma^\mu)^\alpha_\beta \partial_\mu]_{\alpha, \beta=0}^3$ where γ^μ are the well known Dirac γ -matrices [11, 12].

We call an $n \times n$, $n \in \mathbb{N}$ matrix $B = [b^\alpha_\beta]_{\alpha, \beta=1}^n$ a (linear) matrix operator¹⁷ if b^α_β are (linear) operators acting on the space K^1 of C^1 functions $f : M \rightarrow \mathbb{C}$. If $\{f_\nu^0\}$ is a basis in

¹⁷Also a good term for such an object is (linear) *matrixor*.

the set $M(n, 1)$ of $n \times 1$ matrices with the μ -th element of f_ν^0 being $(f_\nu^0)^\mu := \delta_\nu^\mu$,¹⁸ then by definition

$$B\psi := B(\psi) := B \cdot (\psi) := \sum_{\alpha, \beta=1}^n (b_\beta^\alpha (\psi_0^\beta)) f_\alpha^0, \quad \psi = \psi_0^\beta f_\beta^0 \in M(n, 1). \quad (\text{B.1})$$

For instance, we have $\partial\psi = \sum_{\alpha, \beta, \mu=0}^3 (\gamma^\mu)^\alpha_\beta (\partial_\mu \psi_0^\beta) f_\alpha^0$.

To any constant matrix $C = [c_\beta^\alpha]$, $c_\beta^\alpha \in \mathbb{C}$, corresponds a matrix operator $\overline{C} := [c_\beta^\alpha \text{id}_{K^1}]$. Since $C\psi \equiv \overline{C}\psi$ for any ψ , we identify C and $\overline{C}$ and will make no difference between them.

The multiplication of matrix operators, denoted by $\odot$, is a combination of matrix multiplication (denoted by $\cdot$) and maps (operators) composition (denoted by $\circ$). If $A = [a_\beta^\alpha]$ and $B = [b_\beta^\alpha]$ are matrix operators, the product of A and B is also a matrix operator such that

$$AB := A \odot B := \left[\sum_\mu a_\mu^\alpha \circ b_\beta^\mu \right]. \quad (\text{B.2})$$

One can easily show that this is an associative operation. It is linear in the first argument and, if its first argument is linear matrix operator, then it is linear in its second argument too. For constant matrices (see above), the multiplication (B.2) coincides with the usual matrix one.

Let $\{f_\alpha(x)\}$ be a basis in $M(n, 1)$ depending on $x \in M$ and $f(x) := [f_\alpha^\beta(x)]$ be defined by the expansion $f_\alpha(x) = f_\alpha^\beta(x) f_\beta^0$. The *matrix of a matrix operator* $B = [b_\beta^\alpha]$ with respect to $\{f_\alpha\}$, i.e. the matrix of the matrix elements of B considered as an operator, is also a matrix operator $\mathbf{B} := [B_\beta^\alpha]$ such that

$$B\psi|_x =: \sum_{\alpha, \beta} (B_\beta^\alpha(\psi^\beta))|_x f_\alpha(x), \quad \psi(x) = \psi^\beta(x) f_\beta(x). \quad (\text{B.3})$$

Therefore

$$\varphi = B\psi \iff \varphi = \mathbf{B}\psi, \quad (\text{B.4})$$

where $\psi \in M(n, 1)$ is the matrix of the components of ψ in the basis given. Comparing (B.1) and (B.3), we get

$$B_\beta^\alpha = \sum_{\mu, \nu} (f^{-1}(x))_\mu^\alpha b_\nu^\mu \circ (f_\beta^\nu(x) \text{id}_{K^1}). \quad (\text{B.5})$$

For any matrix $C = [c_\beta^\alpha] \in GL(n, \mathbb{C})$, considered as a matrix operator (see above), we have

$$\mathbf{C} = [C_\beta^\alpha(x)], \quad C_\beta^\alpha(x) = (f^{-1}(x))_\mu^\alpha c_\nu^\mu f_\beta^\nu, \quad C(f_\beta) = C_\beta^\alpha f_\alpha. \quad (\text{B.6})$$

Therefore, as one can expect, the matrix of the unit matrix $\mathbb{1}_n = [\delta_\alpha^\beta]_{\alpha, \beta=1}^n$ is exactly the unit matrix,

$$\mathbf{1}_n = \mathbb{1}_n. \quad (\text{B.7})$$

¹⁸ $\delta_\nu^\mu = 0$ for $\mu \neq \nu$ and $\delta_\nu^\mu = 1$ for $\mu = \nu$. From here to equation (B.10) in this appendix the Greek indices run from 1 to $n \geq 1$.

If $\{f_\mu(x)\}$ does not depend on x , i.e. if f_α^β are complex constants, and b^μ_ν are linear, than (B.5) implies

$$B^\alpha_\beta = (f^{-1}(x))^\alpha_\mu f^\mu_\beta(x) b^\mu_\nu \quad \text{for } \partial f^\beta_\alpha(x)/\partial x^\mu = 0. \quad (\text{B.8})$$

In particular, for a linear matrix operator B , we have

$$\mathbf{B} = B \quad \text{for } f^\alpha_\beta = \delta^\alpha_\beta \quad (\text{i.e. for } f_\alpha(x) = f^0_\alpha). \quad (\text{B.9})$$

Combining (B.2) and (B.5), we deduce that the matrix of a product of matrix operators is the product of the corresponding matrices:

$$C = A \odot B \quad \Longleftrightarrow \quad \mathbf{C} = \mathbf{A}\mathbf{B} = \mathbf{A} \odot \mathbf{B}. \quad (\text{B.10})$$

After a simple calculation, we find the matrices of $\mathbb{1}_4 \partial_\mu$ and $\not{\partial} = \gamma^\mu \partial_\mu$.¹⁹

$$\partial_\mu = \mathbb{1}_4 \partial_\mu + \mathcal{E}_\mu(x), \quad \not{\partial} = \mathcal{G}^\mu(x) (\mathbb{1}_4 \partial_\mu + \mathcal{E}_\mu(x)) = \mathcal{G}^\mu(x) \partial_\mu \quad (\text{B.11})$$

where $\mathcal{G}^\mu(x) = [\mathcal{G}^{\lambda\mu}_\alpha(x)]^3_{\alpha,\lambda=0}$ and $\mathcal{E}_\mu(x) = [\mathcal{E}^{\lambda}_\mu(x)]^3_{\alpha,\lambda=0}$ are defined via the expansions

$$\gamma^\mu f_\nu(x) =: \mathcal{G}^{\lambda\mu}_\nu(x) f_\lambda(x), \quad \partial_\mu f_\nu(x) =: \mathcal{E}^{\lambda}_{\mu\nu}(x) f_\lambda(x). \quad (\text{B.12})$$

So, $\mathcal{G}^\mu$ is the matrix of γ^μ considered as a matrix operator (see (B.6)).²⁰ Evidently

$$\not{\partial} = \not{\partial} \quad \text{for } f_\alpha(x) = f^0_\alpha. \quad (\text{B.13})$$

Combining (B.6), (B.7), and (B.11), we get that the matrix of the matrix operator $i\hbar \not{D} - mc\mathbb{1}_4$, entering in the Dirac equation (4.1), is

$$i\hbar \not{D} - mc\mathbb{1}_4 = i\hbar \mathcal{G}^\mu(x) (\mathbb{1}_4 \partial_\mu + \mathcal{E}_\mu(x)) - mc\mathbb{1}_4, \quad (\text{B.14})$$

so that

$$i\hbar \not{D} - mc\mathbb{1}_4 = i\hbar \not{D} - mc\mathbb{1}_4 \quad \text{for } f_\alpha(x) = f^0_\alpha. \quad (\text{B.15})$$

References

- [1] Bozhidar Z. Iliev. Quantum mechanics from a geometric-observer's viewpoint. *Journal of Physics A: Mathematical and General*, 31(4):1297–1305, January 1998.
LANL arXiv server, E-print No. quant-ph/9803083, 1998.
- [2] Bozhidar Z. Iliev. Fibre bundle formulation of nonrelativistic quantum mechanics. I. Introduction. The evolution transport. *Journal of Physics A: Mathematical and General*, 34(23):4887–4918, 2001.
LANL arXiv server, E-print No. quant-ph/9803084, 1998.
- [3] Bozhidar Z. Iliev. Fibre bundle formulation of nonrelativistic quantum mechanics. II. Equations of motion and observables. *Journal of Physics A: Mathematical and General*, 34(23):4919–4934, 2001.
LANL arXiv server, E-print No. quant-ph/9804062, 1998.

¹⁹Here and below the Greek indices run from 0 to 3.

²⁰We denote the matrix of γ^μ by $\mathcal{G}^\mu$ instead of $\gamma^\mu(x)$ as usually [11, 12] by γ is denoted the matrix 3-vector $(\gamma^1, \gamma^2, \gamma^3)$.

- [4] Bozhidar Z. Iliev. Fibre bundle formulation of nonrelativistic quantum mechanics. III. Pictures and integrals of motion. *Journal of Physics A: Mathematical and General*, 34(23):4935–4950, 2001.
LANL arXiv server, E-print No. quant-ph/9806046, 1998.
- [5] Bozhidar Z. Iliev. Fibre bundle formulation of nonrelativistic quantum mechanics. IV. Mixed states and evolution transport's curvature. *International Journal of Modern Physics A*, 17(2):229–243, 2002.
LANL arXiv server, E-print No. quant-ph/9901039, 1999.
- [6] Bozhidar Z. Iliev. Fibre bundle formulation of nonrelativistic quantum mechanics. V. interpretation, summary, and discussion. *International Journal of Modern Physics A*, 17(2):245–258, 2002.
LANL arXiv server, E-print No. quant-ph/9902068, 1999.
- [7] Bozhidar Z. Iliev. Fibre bundle formulation of nonrelativistic quantum mechanics (full version).
LANL arXiv server, E-print No. quant-ph/0004041, 2000, April 2000.
- [8] E. Prugovečki. *Quantum mechanics in Hilbert space*, volume 92 of *Pure and applied mathematics*. Academic Press, New York-London, second edition, 1981.
- [9] Bozhidar Z. Iliev. Linear transports along paths in vector bundles. I. General theory. JINR Communication E5-93-239, Dubna, 1993.
- [10] Bozhidar Z. Iliev. Normal frames and linear transports along paths in vector bundles.
LANL arXiv server, E-print No. gr-qc/9809084, 1998.
- [11] C. Itzykson and J.-B. Zuber. *Quantum field theory*. McGraw-Hill Book Company, New York, 1980. Russian translation (in two volumes): Mir, Moscow, 1984.
- [12] J. D. Bjorken and S. D. Drell. *Relativistic quantum mechanics*, volume 1. McGraw-Hill Book Company, New York, 1964. Russian translation: Nauka, Moscow, 1978.
- [13] A. M. L. Messiah. *Quantum mechanics*, volume II. North Holland, Amsterdam, 1962. Russian translation: Nauka, Moscow, 1979.
- [14] N. N. Bogolyubov and D. V. Shirkov. *Introduction to the theory of quantized fields*. Nauka, Moscow, third edition, 1976. In Russian. English translation: Wiley, New York, 1980.
- [15] N. F. Nelipa. *Physics of elementary particles*. Vŷshaya shkola, Moscow, 1977. (In Russian).
- [16] A. M. L. Messiah. *Quantum mechanics*, volume I. North Holland, Amsterdam, 1961. Russian translation: Nauka, Moscow, 1978.
- [17] Bozhidar Z. Iliev. Linear transports along paths in vector bundles. II. Some applications. JINR Communication E5-93-260, Dubna, 1993.
- [18] *Physical encyclopedia*, chief editor Prohorov A. M., volume 1, Moscow, 1988. Sovetskaya Entsiklopediya (Soviet encyclopedia). (In Russian).
- [19] Bozhidar Z. Iliev. Transports along maps in fibre bundles. JINR Communication E5-97-2, Dubna, 1997.
LANL arXiv server, E-print No. dg-ga/9709016, 1997.

- [20] Bozhidar Z. Iliev. Linear transports along paths in vector bundles. V. Properties of curvature and torsion. JINR Communication E5-97-1, Dubna, 1997.
LANL arXiv server, E-print No. dg-ga/9709017, 1997.